

Graveley Lab CRISPR Genome Editing followed by RNA-seq Biosample Preparation and Characterization Document

Wet Lab: Sara Olson and Lijun Zhan

Computational Lab: Xintao Wei

PI: Brenton Graveley

Department of Genetics and Genome Sciences

UConn Institute for Systems Genomics

UConn Health

400 Farmington Avenue

Farmington, CT 06030 USA

October 13, 2020

Project: ENCORE

Grant: U41HG009889

Sample Description: CRISPR genome editing of DHX57 in K562 cells

Cell Line: K562

RNA ID: DHX57-BGKcLV17-12

ENCODE BIOSAMPLE ACCESSION: ENCBS317WWB

This document contains the protocols used to clone target guide sequence, to generate lentiCRISPRv2-gRNA particles, transduction of K562 cells, harvesting of RNA, characterization of the RNA integrity and measurement of target knockdown efficiency by both qRT-PCR and Western blotting.

Target guide sequence cloning protocol - Adopted from ZhangLab's

In this portion of the protocol we will clone the target sequence into the lentiCRISPRv2.

Item	Info
Target	DHX57-human
gRNA ID	BGC0000440
gRNA sequence	CAGACTGTTATACTGCACCA

In order to clone the target sequence into the lentiCRISPRv2, synthesize two oligos of the following form. All plasmids have the same overhangs after BsmBI digestion.



Lentiviral vector digestion, oligo annealing and cloning into digested vector

1. Digest and dephosphorylate 5ug of the lentiviral CRISPR plasmid with BsmBI for 30 min at 37°C:

Reagent	Quantity
lentiCRISPRv2	5 μ g
FastDigest BsmBI (Fermentas)	3 μ l
FastAP (Fermentas)	3 μ l
10X FastDigest Buffer	6 μ l
100 mM DTT	0.6 μ l
ddH ₂ O (freshly prepared)	X μ l
Total volume	60 μ l

2. Gel purify digested plasmid using QIAquick Gel Extraction Kit and elute in EB.

If BsmBI digested, a ~2kb filler piece should be present on the gel. Only gel purify the larger band. Leave the 2kb band.

3. Phosphorylate and anneal each pair of oligos:

Reagent	Quantity
Oligo 1 (100 μ M)	1 μ l
Oligo 2 (100 μ M)	1 μ l
10X T4 Ligation Buffer (NEB)	1 μ l
T4 PNK (NEB M0201S)	0.5 μ l
ddH ₂ O	6.5 μ l
Total volume	10 μ l

Please use the T4 Ligation Buffer since the buffer supplied with the T4 PNK enzyme does not include ATP (or supplement to 1mM ATP).

Put the phosphorylation/annealing reaction in a thermocycler using the following parameters:

37 °C 30 min
95 °C 5 min and then ramp down to 25 °C at 5 °C/min

4. Dilute annealed oligos from Step 3 at a 1:200 dilution into sterile water or EB.
5. Set up ligation reaction and incubate at room temperature for 10 min:

Reagent	Quantity
BsmBI digested plasmid from Step 2 (50 μ g)	X μ l
diluted oligo duplex from Step 4	1 μ l
2X Quick Ligase Buffer (NEB)	5 μ l
ddH ₂ O	X μ l
Subtotal volume	10 μ l
Quick Ligase	1 μ l
Total volume	11 μ l

Also perform a negative control ligation (vector-only with water in place of oligos) and transformation.

6. Transformation into Stbl3 bacteria.

Lentiviral transfer plasmids contain Long-Terminal Repeats (LTRs) and must be transformed into recombination-deficient bacteria. We use home-made Stbl3 (propagated from Invitrogen C7373-03) and get excellent plasmid yields. Although other RecA-strains may work, we have found the most consistent transformations and yields using Stbl3.

Protocol for producing lentiCRISPRV2-gRNA particles

Day 1

1. Plate 0.8-1x10⁶ 293 T cells (catalog number: CRL-11268, ATCC) in each well of 6-well plate with 10 % FBS (catalog number: 30-2020, ATCC) DMEM (catalog number: 11995-065, Life technologies) medium without penicillin and streptomycin.
2. Incubate overnight. Cells should be 70-80% confluent.

Day 2

1. In polypropylene tubes, make a cocktail for each transfection as follows:

Reagent	Quantity
lentiCRISPRv2-gRNA	1 μ g
psPAX2 Packaging DNA	750 ng
pCMV-VSV-G DNA	250 ng
serum-free OPTI-MEM	to 100 μ l

2. Add 6 μ l of FuGENE HD Transfection reagent (Catalog number: E2311, Promega) to the tube (FuGENE:DNA=3:1)
3. Incubate for 20 minutes at room temperature.
4. Gently add the DNA mix dropwise to cells.
5. Incubate the cells at 37 °C for 12-15 hr.

Day 3

1. In the morning, change the media to remove the transfection reagent, wash with PBS once and add 1.5 ml fresh media +10% FBS + penicillin/streptomycin.

Day 4

1. Harvest media from cells, store at 4 °C.
2. Add 1.5 ml fresh media.

Day 5

1. Harvest the media from the cells and pool with the media collected on Day 4.
2. Spin the media at 1250 rpm for 5 min to remove cells.
3. Freeze the virus stock at -40 °C.

Lentiviral CRISPR Protocol

For K562 cells

Source: ATCC CCL-243 (lot 59300853)

Growth Media for K562 cells:

500 ml RPMI-1640 with glutamine medium (Hyclone, SH30027.01)
50 ml Fetal Bovine Serum (FBS) (10% Final Concentration) (Hyclone, SH30071.03)
5 ml Pen-Strep (1% Final Concentration) (Invitrogen, 15140-163)

Culturing

1. Remove a frozen stock vial of K562 cells from liquid nitrogen.
2. Thaw it in a 37 °C water bath.
3. As soon as it thawed, transfer the cells into the growth medium.
4. Centrifuge at 1000 rpm for 5 minutes.
5. Resuspend the cells in the growth medium such that the final cell density is about 1×10^5 cells/ml.
6. Split the cells every 2 to 3 days.
7. Grow the cells to required numbers.

Prepare cells for transduction

Day 0

1. Plate 0.8-1ml of lentiCRISPRv2-gRNA particles in each well of 12-well plates.
2. Add $5-7 \times 10^5$ cells to appropriate wells with 8 $\mu\text{g}/\text{ml}$ of polybrene (Catalog Number H9268, Sigma-Aldrich), incubate at 37 °C for 2 hours, and then add 1 ml of complete medium.

Day 1

1. After 24 hrs, change to fresh media (2 ml) with 3 $\mu\text{g}/\text{ml}$ of puromycin.

Day 3

1. Change to fresh media (2 ml) with 3 $\mu\text{g}/\text{ml}$ of puromycin.

Day 5

1. Change to fresh media (2 ml) with 3 $\mu\text{g}/\text{ml}$ of puromycin.

Day 6

1. Harvest half of the cells to prepare RNA and half of the cells to prepare a protein lysate for western blotting.

RNA Isolation

RNA isolation is performed using a Promega Maxwell®16 Instrument and the Maxwell®16 LEV simplyRNA Cells Kits (Catalog Number AS1270).

1. Pellet cells at 300 x g for 3 minutes and remove medium.

2. Add 200 μl of chilled 1-Thioglycerol/Homogenization solution to the cell pellet and vortex until the pellet is dispersed.
3. Add 200 μl of lysis buffer and vortex vigorously for 15 sec to mix.
4. Transfer all 400 μl lysate to well 1 of the Maxwell 16 LEV cartridge.
5. Add 5 μl of DNase I solution to well 4 of the cartridge.
6. Put elution tubes with 40-50 μl of nuclease-free water and LEV plungers in the cartridge.
7. Transfer the Maxwell 16 LEV cartridge rack containing prepared cartridges on the Maxwell 16 Instrument.
8. Push Run/Stop button to start run.

RNA Quality Control

The quality of the RNA is measured using an Agilent TapeStation Instrument with the RNA screen tape (Catlog Number 5067-5576).




Figure 1: Agilent TapeStation image of 1 μl of total RNA sample.

qRT-PCR Assay to Monitor mRNA Target Knock-down Efficiency

cDNA Synthesis

This assay uses the iScript cDNA Synthesis Kit from BIO-RAD (Catalog number: 170-8891)

1. Reaction Setup:

Reagent	Quantity
5x iScript reaction mix	2 μ l
iScript reverse transcriptase	0.5 μ l
Nuclease-free water	x μ l
RNA template (200 ng)	x μ l
Total volume	10 μ l

2. Reaction Protocol:

Time	Temperature
5 minutes	25 °C
30 minutes	42 °C
5 minutes	85 °C
Hold	4 °C

qPCR Assay

This assay uses Phusion High-Fidelity DNA Polymerase from NEB (Catalog number: M0530L) and SYBR Green from Invitrogen (Catalog number: S7563)

1. Reaction setup:

Reagent	Quantity
5X Phusion HF Buffer	4 μ l
10 mM dNTPs	0.4 μ l
10 μ M Forward Primer	1 μ l
10 μ M Reverse Primer	1 μ l
Template (1:20 of cDNA reaction)	1 μ l
Phusion DNA Polymerase	0.2 μ l
SYBR Green (10,000 X)	0.1 μ l
Nuclease-free water	to 20 μ l
Total volume	20 μ l

2. Reaction Protocol:

STEP	TEMP	TIME
Initial Denaturation	98 °C	30 seconds
35 Cycles	98 °C	10 seconds
	58 – 66 °C	15 seconds

3. Data Analysis:

Data analysis is performed using the $2^{-\Delta\Delta C_t}$ Method.




Figure 2: qRT-PCR analysis of depletion level of the target RNA binding protein in control and knockdown cells. The percent depletion was calculated in the RNA sample isolated from K562s transduced with a gRNA targeting DHX57 (ENCODE Biosample ENCSR359ICP and ENCBS317WWB) in comparison to K562 cells transduced with a control non-target gRNA (ENCODE Biosample ENCSR516TLX and ENCBS813SYB). The efficiency of depletion is normalized using GAPDH as a control.

Western Blot Assay to Monitor Protein Target Knockdown Efficiency

A western blot is performed to determine the knockdown efficiency of the target RNA binding protein. For this biosample, the following antibodies were used:

RNA binding protein primary antibody: DHX57

Loading control primary antibody: GAPDH

Wes Protocol

12-230 kDa Master kit with split Running Buffer from Proteinsimple

A. PREPARE STANDARD PACK REAGENTS

1. Add 40 μL deionized water to make a 400 mM solution of the DTT.
2. Add 20 μL 10X Sample buffer and 20 μL prepared 400 mM DTT to make 5X Fluorescent Master Mix.
3. Add 16 μL deionized water, 2 μL 10X Sample Buffer and 2 μL prepared 400 mM DTT solution to make Biotinylated Ladder.

B. PREPARE YOUR SAMPLES

1. Combine 1 part 5X Fluorescent Master Mix with 4 parts lysate in a microcentrifuge tube (If needed, dilute the lysate with 0.1X Sample Buffer).
2. Denature the samples and biotinylated ladder at 70 °C for 10 min.

C. PREPARE REAGENTS FROM DETECTION MODULE

1. Dilute primary antibody with antibody diluent 2.
2. The supplied secondary antibody is ready to use without dilution.
3. Combine 150 μL Luminol-S and 150 μL Peroxid in a microcentrifuge tube. Gently pipette up and down to mix and store on ice.

D. PIPETTE YOUR PLATE (IMMUNOASSAY)



For more consistent results, keep the lid on between reagent additions. Centrifuge the plate for 5 minutes at 2500 rpm at room temperature.



E. START WES

1. Load the desired assay in Compass software v2.7 or higher.
2. Open Wes' door.
3. Insert a capillary cartridge into the cartridge holder. The interior light will change from orange to blue.
4. Remove the assay plate lid. Hold plate firmly on bench and carefully peel off evaporation seal. Pop any bubbles observed in the Separation Matrix wells with a pipette tip.
5. Place the assay plate on the plate holder
6. Close Wes' door.
7. Click the Start button in compass.
8. When the run is complete, discard the plate and cartridge.




Figure 3: Western Blot Analysis of depletion level of the target RNA binding protein in control and knockdown cells. Lane 1: Molecular weight marker. Lane 2: 30 μg of protein from K562 transduced with a control, non-target gRNA (ENCODE Biosample ENCSR516TLX and ENCBS813SYB). Lane 3: 30 μg of protein from K562 transduced with a gRNA targeting DHX57 (ENCODE Biosample ENCSR359ICP). Lane 4: 30 μg of protein from K562 transduced with a gRNA targeting DHX57 (ENCODE Biosample ENCBS317WWB). Samples were separated by SDS-PAGE, transferred to a membrane and blotted using antibodies against DHX57 (ENCODE Antibody ENCAB163YWK) and GAPDH as controls.